

DNA Methylation and Primary Immune Thrombocytopenia

Huiyuan Li, Min Xuan and Renchi Yang

DNA methylation is a heritable, stable, and also reversible way of DNA modification; it can regulate gene expression without changing the nucleotide sequences. Because it takes part in regulation of immune responses, the loss of methylation homeostasis in immune cells will result in autoimmune disease by inducing aberrant gene expression. Primary immune thrombocytopenia (ITP) is an acquired autoimmune disease with many immune deficiencies. Recently, it was well documented that abnormal DNA methylation is also involved in the etiology of ITP. In this review, we elucidate the role of DNA methylation in autoimmune diseases by summarizing the DNA methylation-sensitive genes and the relationship between DNA methylation and ITP.

Semin Hematol 50:S116–S126. © 2013 Published by Elsevier Inc.

Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

The autoimmune diseases are a series of disorders with various and complex defects in the immune system; the etiologies for some of them remain unclear today. Although DNA sequence changes have been claimed to be associated with susceptibility to disease onset, the different concordance rate of autoimmune diseases between monozygotic twins indicates that environment also participates in the development of autoimmunity.^{1–4} These environmental factors effect immunity mainly by altering epigenetic regulation, which ultimately results in changes in gene expression.^{5,6} DNA methylation is an epigenetic process that refers to heritable chromatin-based mechanisms in the regulation of gene expression without DNA alternation. This modification of DNA is time- and cell-specific, and thus it plays an important role in the regulation of many pathology processes, such as embryonic development, X chromosome inactivation, genomic imprinting, and cellular differentiation.^{7–10}

DNA METHYLATION

DNA methylation is mediated by DNA methyltransferases (DNMTs) through the addition of a methyl group from S-adenosylmethionine (SAM) to the fifth carbon of cytosine residues in CG dinucleotides.¹¹ There are five members in DNMT group: DNMT1, DNMT2, DNMT3A, DNMT3B, and DNMT3L. The function of DNMT1 is to methylate the hemimethylated sites generated during DNA replication. As the maintenance DNMT, it is implicated in stabilizing DNA methylation patterns during cell division.¹² DNMT3A and 3B are known as de novo DNA methyltransferases; they are responsible for the establishment of DNA methylation patterns during fetal development. In addition, DNMT3A and DNMT3B also exhibit non-overlapping functions in this procedure, with DNMT3B being specifically required for methylation of centromeric minor satellite repeats.^{13,14} DNMT2 is not essential for global de novo or maintenance methylation of DNA because of its weak DNA methyltransferase activity; instead, it acts on methylation of small RNA.^{15–17} DNMT3L is crucial for the establishment of maternal genomic imprints but lacks key methyltransferase motifs; it is more likely to act as a regulator of methylation by recruiting histone deacetylase (HADC) and stimulating de novo methylation by DNMT3A.^{18–20}

DNA methylation is a stable but reversible epigenetic modification; thus DNA demethylation, which contributes to changes in the DNA methylation pattern, is also vital for biological processes. DNA demethylation includes passive and active mechanisms. Passive mechanisms require DNA replication

State Key Laboratory of Experimental Hematology, Institute of Hematology & Blood Diseases Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, PR China. Publication of this article was supported by the International Cooperative ITP Study Group (ICIS).

Conflicts of interest: none.

Address correspondence to Professor Renchi Yang, State Key Laboratory of Experimental Hematology, Institute of Hematology and Blood Diseases Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, 288 Nanjing Rd, Tianjin 300020, PR China. E-mail: rcyang65@yahoo.com

0037-1963

© 2013 Published by Elsevier Inc. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).
<http://dx.doi.org/10.1053/j.seminhematol.2013.03.022>




Figure 1. DNA methylation can repress gene transcriptions. DNA methylation is mediated by DNA methyltransferases (DNMTs) through the addition of a methyl group from S-adenosylmethionine (SAM) to the fifth carbon of cytosine residues in CG dinucleotide. On the one hand, these methyl groups can inhibit gene expression directly by blocking transcription complex binding to DNA. On the other hand, methylated DNA may be bound by proteins known as methyl-CpG-binding domain proteins (MBDs). MBDs can further recruit histone deacetylases (HADC), that eventually alter chromatin structures and lead to inaccessible transcription. Red hexagon stands for methyl group. Pol II: RNA polymerase II; TF: transcription factors.

by inhibition of DNMTs.²¹ The active process is independent of DNA replication; it occurs by removing the methyl group by demethyltransferases, such as methyl-CpG-binding domain (MBD) proteins.²²

DNA methylation can repress transcription through several mechanisms (Figure 1). First, methyl groups inhibit gene expression directly by blocking transcription complex binding to DNA.²³ Second, MBDs (MBD1, MBD2, MBD3, MBD4, and MECP2) are attracted to the methylated CpG dinucleotide. Since the binding of these proteins on methylated DNA also prevented the binding of transcription complex, MBDs were reported to be transcriptional repressors as well as DNA demethylases. In addition, MBDs can recruit HADC, eventually altering chromatin structures and leading to inaccessible transcription.^{24,25}

The DNA methylated status of a genome is influenced by several factors. First, the 5-methylcytosine content in DNA declines with age, suggesting that the methylated level is related to the proliferative potential of organs.²⁶ Meanwhile, the DNA methylated level is different between males and females. As women have two X chromosomes, a high level of DNA methylation is used to inactivate one X chromosome. This X-inactivation results in women having equivalent X chromosome gene products as males.⁹ In addition, food, drug, and environmental factors also impact DNA methylation.^{5,6,27,28}

DNA METHYLATION AND AUTOIMMUNITY

Normal T cells treated with the DNA methylation inhibitor 5-azacytidine induce autologous B-cell differentiation and T-cell proliferation, and ultimately

initiate autoimmunity.^{29,30} Thus, DNA methyltransferase inhibitors, such as procainamide and 5-azacytidine, are sufficient to induce a lupus-like disease.³¹ In addition, the lymphocytes of systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA) manifest global DNA hypomethylated status due to the abnormal expressions of DNMTs and MBDs,³² which further demonstrates that DNA methylation plays a key role in autoimmune diseases.

What is the reason for hypomethylation in some autoimmune diseases? There is evidence that the impaired ERK and ras-MAPK pathway signaling in T cells results in decreased expression of DNMT1 and DNA hypomethylation.^{33,34} This defective methylation capacity finally generates autoimmunity mainly through inducing overexpression of methylation-sensitive genes that are engaged in autoreactive immune responses in lymphocyte, including *IL-4*, *IL-6*, *CD70*, and *FoxP3* among others (Table 1).

CD11a

Lymphocyte function-associated antigen 1 (LFA-1, CD11a/CD18) is a leukocyte cell surface heterodimer. It promotes intercellular adhesion by binding to members of the intercellular adhesion molecule family in immunological and inflammatory reactions.³⁵ The expression of LFA-1 is tissue-specific through the regulation of methylation and chromatin structure.³⁶ Normal T cells treated with DNA methylation inhibitors had increased CD11a expression directly for demethylation of the CD11a promoter.^{37,38} Lu et al. found that the specific sequence flanking the CD11a promoter was hypomethylation in T cells from patients with active lupus or in normal T cells treated with 5-azacytidine and procainamide. Patch methylation of this region can suppress CD11a promoter function, which illustrated that altered methylation of specific genes may be relevant to the pathogenesis of lupus.³⁹

CD70

CD70 is a costimulatory molecule of the tumor necrosis factor super family. It is expressed on activated dendritic cells, B cells, and T cells. CD27, the CD70 receptor, is also expressed on activated immunocytes. CD70-CD27 signaling provides a costimulatory signal for T-cell activation and survival. CD70 is another methylation-sensitive gene that is overexpressed in normal T cells treated with DNA methylation inhibitors, and these overexpressed CD70 contribute to overstimulated B cells and increased IgG production.^{40,41} In SLE patients, CD70 expression is significantly elevated and

Table 1. Summary of DNA Methylation Involved in Autoimmune Diseases

Disease	Global DNA Methylation Levels	Methylation-Associated Enzymes	Methylation-Sensitive Genes		
			Genes	Methylation Status	Expression
SLE	Hypomethylation ^{79,80}	DNMT1 ↓ ⁷⁹ DNMT3A, 3B ↓ ⁸¹ MBD2, MeCP ↑ ⁸²	CD70, ⁴² CD40, ⁷³ IL-4 and IL-6, ⁵² IL-13, ⁸³ IL-2, ⁶⁹ perforin, ⁶⁰ CD11a, ³⁹ CD5, ⁷⁷ protein phosphatase 2, ⁸⁵ IL-10 and IL-1 receptor 2, ³² 5-hydroxytryptamine receptor 1A, ⁸⁴ others ^{86–89}	Hypomethylation	Increase
RA	Hypomethylation ⁹⁰	DNMT1 ↑ MBD2 ↑ ⁹⁰	CD40L, ⁷¹ IL-6, ⁶³ CXCL12, ⁹¹ IL-1 receptor 2 ³² IL-10 ^{32,92}	Hypomethylation	Increase
Psoriasis	Inconsistent ^{94,95}	DNMT1 ↑ NBD2, MeCP ↓ ⁹⁵	Death receptor-3 ⁹³ p15 and p21, ⁹⁶ protein tyrosine phosphatase-1 ⁹⁸ p14 ⁹⁵	Inconsistent Hypermethylation Hypomethylation	Decrease Decrease Increase
MS	Hypomethylation ¹⁰⁰		p16 ^{97,99} Protein tyrosine phosphatase-1 ¹⁰¹ Peptidyl argininedeiminase 2 ^{100,102}	Inconsistent Hypermethylation	Decrease Decrease
SS	Hypomethylation ^{72,79,103}	DNMT1 ↓ MBD3, MBD4 ↓ ⁷⁹	CD70, ^{44,99} CD40L ⁷²	Hypomethylation	Increase
T1D	Hypermethylated ¹⁰⁴	DNMT3B ↑ ⁵⁶	Insulin, ¹⁰⁴ FoxP3 ⁵⁶	Hypermethylation	Decrease
UC	Hypermethylated	DNMT1, DNMT3B ↑ ¹⁰⁵	Multidrug resistance 1, ¹⁰⁶ p16, ¹⁰⁷ p14, ¹⁰⁸ estrogen receptor-1, tumor suppressor candidate-3, ¹⁰⁹ others ¹⁰⁵ Protease-activated receptor 2 ¹¹⁰	Hypermethylation Hypomethylation	Decrease Increase

Abbreviations: SLE, systemic lupus erythematosus; RA, rheumatoid arthritis; MS, multiple sclerosis; SS, systemic sclerosis; T1D, type 1 diabetes; UC, ulcerative colitis; DNMT, DNA methyltransferase; MBD, methyl-CpG-binding domain protein.

correlates with a decrease in CD70 promoter methylation in CD4⁺ T cells compared to healthy controls, as the consequence of the decreased expression of DNMT1.⁴² Besides SLE, the reduction of methylation in CD70 promoter also causes overexpression of CD70 in many other autoimmune diseases, such as primary Sjögren's syndrome, systemic sclerosis (SS), and immune thrombocytopenia (ITP).^{43–45}

Interleukin-4 and Interferon-γ

Interferon (IFN)-γ is the typical cytokine of Th1 cells, which is mainly engaged in macrophage activation. Interleukin (IL)-4 is the typical cytokine of Th2 cells, which generally respond by inducing antibody production. The balance between Th1 and Th2 subsets has been implicated in the regulation of many immune responses.⁴⁶ CD4⁺ T cells

treated with 5-azacytidine or procainamide secrete relatively large amounts of IL-4 and IFN- γ ,^{31,47} implying that DNA methylation participates in T-cell differentiation. Further studies show that the *IFN- γ* and *IL-4* genes are methylated in human naive CD4⁺ T cells. After stimulation via T-cell receptor (TCR), the IFN- γ promoter becomes hypermethylated in Th2 cells, whereas it is hypomethylated in Th1 cells. Hypermethylation in Th2 cells results in chromatin condensation and exclusion of cAMP response element binding protein (CREB) from IFN- γ promoter.^{48,49} Accordingly, the promoter of IL-4 locus becomes specifically demethylated in Th2 cells, but becomes methylated during Th1 differentiation.⁵⁰ The hypomethylation of the IL-4 promoter in the pathogenesis of SLE and asthma might be important through upregulating the level of IL-4.^{51,52}

FoxP3

The expression of forkhead box P3 (FoxP3) is regulated by epigenetics. It is unmethylated in natural regulatory T cells (nTreg), but heavily methylated in naive CD4⁺ T cells, activated CD4⁺ T cells, and peripheral transforming growth factor (TGF)- β -induced Tregs. Thus, FoxP3 DNA demethylation constitutes the most reliable criterion for natural Tregs available at present.^{53,54} During T-cell activation in vitro, a DNA demethylation agent, 5-Aza-2'-deoxycytidine, can induce Foxp3 expression in CD4⁺CD25⁺Foxp3⁻ cells via altering methylation status of a conserved element in the 5'-untranslated region of the *Foxp3* gene. In addition, the methylation status within the Foxp3 locus in CD4⁺ T cells is associated with the expression of DNMT1 and DNMT3B, and knocking down the DNMT1 induces FoxP3 expression.^{55,56} In latent autoimmune diabetes (LADA) in adults, the FoxP3 promoter region is hypermethylated in CD4⁺ T cells from LADA patients compared with the healthy cohort, which may contribute to disease onset and progression of LADA.⁵⁶

Perforin

Perforin is a cytotoxic effector molecule expressed in natural killer cells and a subset of T cells, and it contributes to target cell killing.⁵⁷ As a methylation-sensitive gene, the perforin promoter region is hypomethylated in primary CD8⁺ cells, which express perforin, but is largely methylated in primary CD4⁺ T cells, which do not.^{58,59} It has been reported that perforin is overexpressed in CD4⁺ T cells from active, but not inactive SLE patients, and this overexpression is related to hypomethylation of the perforin promoter in primary CD4⁺ T cells.^{60,61}

IL-6

IL-6 activates various cell types carrying the membrane bound IL-6 receptor; its signaling plays a pivotal role in controlling the differentiation and activation of T and B lymphocytes. Treatment with 5-aza-2'-deoxycytidine induces a high level of IL-6 in MCF-7 (a breast carcinoma cell line) via its gene hypomethylation, which demonstrated that IL-6 expression is influenced by DNA methylation.⁶² In SLE patients, the hypomethylation of the CpG islands of the IL-6 promoter occurred in T cells and correlates with symptom severity.⁵² In addition, the hypomethylated status of a single CpG in the IL-6 promoter region, such as CpG motif at -74 bp or -1099 bp may augment serum levels of IL-6, implicating a potential role in the pathogenesis of RA and chronic periodontitis.^{63,64} In turn, IL-6 may exert many epigenetic changes in cells via increasing expression and activity of DNMTs.⁶⁵

IL-2

IL-2, a key cytokine affecting proliferation and activation of T and B lymphocytes, is also regulated by DNA methylation. During T-cell, B-cell, and macrophage activation, a small region in the promoter-enhancer of IL-2 was demethylated; this demethylated CpG site recruits Oct-1, and induces changes in histone modifications, finally eliciting *IL-2* gene expression.^{66,67} Furthermore, the expression of several potential enzymes/co-enzymes in relation to the DNA demethylation pathways such as DNMT1, DNMT3A, and MBD4 seemed to be associated with immune cell activation by regulating the expression of IL-2.⁶⁸ Downregulated IL-2 expression is a hallmark of SLE T lymphocytes and is linked to overproduction of the transcription regulatory factor cAMP-responsive element modulator (CREM) α , which binds to a CRE site within the IL-2 promoter and results in epigenetic changes in T lymphocytes through DNMT3A-directed DNA hypermethylation and histone deacetylase-1-directed deacetylation.⁶⁹

CD40 Ligand

CD40 ligand (CD40L), a costimulator expressed on CD4⁺ T cells, is a response to B-cell activation and immunoglobulin (Ig) class-switch; it is a methylation-sensitive gene that is encoded by X chromosome. Normal CD4⁺ T cells treated with demethylation agent overexpress CD40L mRNA and induce autologous B-cell activation and plasma cell differentiation. All of these effects can be reversed by anti-CD40L antibody.⁷⁰ In patients with SLE, RA, primary biliary cirrhosis, or SS, demethylation of CD40L regulatory elements on the inactive

X chromosome contributes to CD40L overexpression in CD4⁺ T cells, which may in part explain the female preponderance of autoimmunity.⁷¹⁻⁷⁴

CD5

CD5⁺ B cells, also called B1 cells, play important roles in the production of high-affinity auto-antibodies; thus they participate in autoimmunity. CD5 has two isoforms: E1A and E1B. E1A is the full-length and transmembrane-directed protein, whereas E1B is a truncated isotype that remains in the cytoplasm.⁷⁵ Accordingly, E1A can translocate the Src homology 2 (SH2)-containing tyrosine phosphatase-1 (SHP-1) to the vicinity of the B-cell receptor and raise its threshold, thereby limiting the response of autoreactive B cells. In contrast, the truncated variant E1B stays in the cytoplasm, where it can bind to SHP-1 and prevent its effects.⁷⁶ The expression of E1B is regulated by DNA methylation. In SLE patients, the promoter for the alternative E1B isoform is demethylated in B cells from patients but not in healthy controls. This finally results in elevated intracellular E1B levels.^{77,78}

Others

In addition to the above-mentioned, a large number of methylation-sensitive genes still refer to the pathogenesis of autoimmune diseases (Table 1). Nonetheless, because of the prevalence of CpG islands in genomes, the methylation-sensitive genes related to autoimmune diseases must be more complex than we can imagine. Further research is clearly needed to better understand methylation-sensitive genes in autoimmunity.

DNA METHYLATION AND ITP

Primary ITP is an acquired autoimmune disease characterized by decreased platelet count due to both increased platelet destruction and insufficient platelet production.¹¹¹ More than one mechanism could contribute to ITP, including autoreactive B lymphocytes, Th1/Tc1 polarization, T-cell-mediated platelet lysis, abnormal circulating Treg cells, and so on.¹¹²⁻¹¹⁶ However, until now, the etiology of ITP was incompletely understood. Like in some autoimmune diseases described above, DNA methylation also participates in the pathophysiology of ITP.

Global DNA Methylation Levels in ITP Patients

S-adenosylmethionine (SAM) and S-adenosylhomocysteine (SAH) are the donor and receptor of methyl group in the cellular methyltransferase reaction; thus SAM/SAH can be used as an indicator of cellular methylation status. The SAM/SAH ratio in

plasma directly correlates with that in lymphocytes, and can reflect cellular methylation levels.^{117,118} We measured serum concentrations of SAM and SAH in ITP patients and normal controls using high-performance liquid chromatography, and found that the plasma SAH level was significantly elevated in ITP patients compared with normal controls.⁸¹ In addition, although the SAM/SAH ratio has a tendency to decrease in ITP patients, no significant differences were observed between ITP patients and controls.^{81,119} Because SAH has a higher affinity for the methyltransferase active site than SAM,¹²⁰ elevated SAH can bind to and inhibit the effect of DNMTs; thus we speculated that lymphocyte DNA hypomethylation may exist in ITP through elevated plasma SAH levels. However, upregulated SAH also can be explained by an increase in DNA methylation, suggesting DNA hypermethylation. Thus further study on the DNA methylation status in ITP patients has been done. Chen et al further assessed global methylation by quantifying the methylcytosine concentration of genomic DNA.¹²¹ The result revealed that CD4⁺ T cells had a low DNA methylcytosine content in ITP patients, which prompted the conclusion that the hypomethylation pattern occurred in CD4⁺ T cells of ITP patients. However, the methylation levels seem irrelevant to clinical parameters, including age, sex, clinical course, and platelet counts.¹²¹ Based on these findings, we speculate that the aberrant DNA methylation may take part in the pathogenesis of ITP.

Abnormal Expression of DNA Methylated Enzymes in ITP Patients

The aberrant DNA methylation status is mostly induced by abnormal expression of methylation associated enzymes/co-enzymes. Tao et al found that the mRNA expressions of DNMT3A and DNMT3B were downregulated in peripheral blood mononuclear cells (PBMCs) of ITP patients compared with normal controls. This might be due to the elevated plasma SAH concentrations.⁸¹ Consistent with this result, EL-Shiekh et al found DNMT3A mRNA expression in PBMCs was significantly decreased, while the plasma SAH level was elevated in patients with ITP compared with healthy controls. Thus they speculated that aberrant DNA methylation status, reflected by decreased mRNA expression of DNMT3A and increased plasma SAH level, may engage in the pathogenesis of ITP.¹¹⁹ However, Ma et al came to a contrary conclusion through the measurement of mRNA expressions of DNMTs in CD4⁺ T cells from ITP patients.⁴⁵ They found that DNMT1, DNMT3A, and DNMT3B were all elevated in CD4⁺ T cells from ITP patients, and speculated that DNMTs also work as demethyltransferases as well as methyltransferases

in ITP patients.⁴⁵ In addition, the detection of mRNA expressions of MBDs showed that the mRNA expression of MBD2 and MBD4 were decreased both in PBMCs and CD4⁺ T cells in ITP patients compared with those of normal controls,¹²¹ which is different from the result observed in SLE patients.⁷⁹ The possible explanation for this discrepancy could be the Th1/Th2 polarization in ITP and the fact that IFN- γ expression was restricted by MBD2.¹²²

Methylation-Sensitive Genes in ITP Patients

As a methylation-sensitive gene, aberrant methylation of CD70 is also involved in the etiology of ITP. Firstly, Ma et al detected the expression of CD70 on CD4⁺CD8⁺ T cells and CD19⁺ B cells and found that both the mRNA and protein expressions of CD70 were both increased in CD4⁺ T cells from ITP patients, but there was no statistical difference in CD8⁺ or CD19⁺ cells between ITP patients and healthy controls.¹²³ They further detected the DNA methylation indices by polymerase chain reaction (PCR) amplification and high-resolution melting analysis, the result of which demonstrated that the methylation levels of the promoter region of CD70 in ITP were lower than those found in healthy controls. In addition, they found a negative correlation between methylation indices and CD70 mRNA levels in ITP patients and healthy controls. Thus the elevated expression of CD70 may be caused by the hypomethylated promoter region of CD70.⁴⁵ This upregulated CD70 can facilitate the survival of T and B lymphocytes, apoptosis of platelets, and secretion of IFN- γ , finally accelerating the progress of ITP.¹²³

It is clear that abnormal T-cell immunity including Th1, Th2, Treg, and cytotoxic T lymphocyte (CTL) promotes the development of ITP. As a methylation-sensitive gene, *FoxP3* is the key transcription factor for differentiation of Tregs. Moreover, IFN- γ , IL-4 and perforin are major cytokines secreted by the Th1, Th2, and CTL, respectively. To determine the role of DNA methylation in ITP, Zhao et al¹²⁴ investigated the relationship between the mRNA expressions of the *IFN- γ* , *IL-4*, *FoxP3*, and *perforin* genes and the methylation status of their respective promoters in PBMCs from ITP patients. However, although the methylation rate of CpGs located at -409 of *perforin* and the whole methylation of the *IFN- γ* gene were higher than those of normal controls, they did not find any negative correlation between the mRNA expression of these genes and the methylation status of their promoters.

Single-Nucleotide Polymorphism and DNA Methylation Defect in ITP Patients

Single-nucleotide polymorphism (SNP) is the most common form of human genetic variation, and may

be influence the expression or activity of the protein that was encoded by the mutated gene. This will ultimately result in abnormal biological processes. Several genes in the DNA methylation pathway have been subject to genetic association studies in ITP patients.

Chen et al detected the -149C>T SNP (rs2424913) in the promoter of DNMT3B, but did not find any significantly difference in either genotypes or allelic distribution between ITP patients and normal controls.¹²⁵ This result was confirmed by Shaheen et al.¹²⁶ In addition, Zhao et al found that there was also no significant difference of another DNMT3B SNP -579G>T (rs1569686) in genotype and allele distribution between the ITP patients and normal controls.¹²⁷ Thus it is shown that neither of the two SNPs of DNMT3B can be used as markers to predict the susceptibility to ITP.

MBD4 is different from the other members of the MBD family. It preferentially binds to 5mCpG -TpG mismatches and removes the mismatched thymine or uracil, and thus it is thought to act as a DNA repair enzyme that minimizes mutations at methyl-CpG.¹²⁸ Zhao et al¹²⁹ investigated the association between MBD4 rs140693 polymorphism and the risk for ITP in a Chinese population. However, there was no significant difference in genotype and allele distribution between the ITP patients and the controls. Similar results were observed between the two groups when stratified by age and disease course, including acute childhood, chronic childhood, acute adult, and chronic adult. In conclusion, MBD4 polymorphism may not be a marker to predict the susceptibility to ITP, at least in the Chinese population.¹²⁹

CONCLUSION

In this review, we have shed light on the important role DNA methylation plays in autoimmune diseases, including ITP, by regulating the expression of methylation-sensitive genes. Thus the identification of a methylation-sensitive gene in ITP will provide a clinical marker of diagnosis, disease progression, and therapy. In addition, for the reversibility of DNA methylation, the site-specific methylated modification may be effective for the treatment of active ITP patients. However, the research about DNA methylation in ITP is still in its preliminary stage, and further experiments are needed to explore its role in ITP.

Acknowledgments

This work was supported in part by grants from the National Natural Science Foundation of China (30670900, 81070397), Ministry of Science and Technology (2011ZX09302-007-04), Ministry of Health

(201202017), and Tianjin Municipal Science and Technology Commission (10JCZDJC19700).

REFERENCES

- Jarvinen P, Kaprio J, Makitalo R, Koskenvuo M, Aho K. Systemic lupus erythematosus and related systemic diseases in a nationwide twin cohort: an increased prevalence of disease in MZ twins and concordance of disease features. *J Intern Med*. 1992;231:67-72.
- Kyvik KO, Green A, Beck-Nielsen H. Concordance rates of insulin dependent diabetes mellitus: a population based study of young Danish twins. *BMJ*. 1995;311:913-7.
- Aho K, Koskenvuo M, Tuominen J, Kaprio J. Occurrence of rheumatoid arthritis in a nationwide series of twins. *J Rheumatol*. 1986;13:899-902.
- Mumford CJ, Wood NW, Kellar-Wood H, Thorpe JW, Miller DH, Compston DA. The British Isles survey of multiple sclerosis in twins. *Neurology*. 1994;44:11-5.
- Baccarelli A, Wright RO, Bollati V, et al. Rapid DNA methylation changes after exposure to traffic particles. *Am J Respir Crit Care Med*. 2009;179:572-8.
- Tarantini L, Bonzini M, Apostoli P, et al. Effects of particulate matter on genomic DNA methylation content and iNOS promoter methylation. *Environ Health Perspect*. 2009;117:217-22.
- Niles KM, Chan D, La Salle S, Oakes CC, Trasler JM. Critical period of nonpromoter DNA methylation acquisition during prenatal male germ cell development. *PLoS One*. 2011;6:e24156.
- Liang P, Song F, Ghosh S, et al. Genome-wide survey reveals dynamic widespread tissue-specific changes in DNA methylation during development. *BMC Genomics*. 2011;12:231.
- Mohandas T, Sparkes RS, Shapiro LJ. Reactivation of an inactive human X chromosome: evidence for X inactivation by DNA methylation. *Science*. 1981;211:393-6.
- Bartolomei MS, Tilghman SM. Genomic imprinting in mammals. *Annu Rev Genet*. 1997;31:493-525.
- Branno M. Determination of S-adenosylmethionine and S-adenosylhomocysteine in nuclei isolated from sea urchin embryos during early development. *Boll Soc Ital Biol Sper*. 1980;56:1769-71.
- Riggs AD. X inactivation, differentiation, and DNA methylation. *Cytogenet Cell Genet*. 1975;14:9-25.
- Okano M, Bell DW, Haber DA, Li E. DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell*. 1999;99:247-57.
- Okano M, Xie S, Li E. Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nat Genet*. 1998;19:219-20.
- Okano M, Xie S, Li E. Dnmt2 is not required for de novo and maintenance methylation of viral DNA in embryonic stem cells. *Nucleic Acids Res*. 1998;26:2536-40.
- Hermann A, Schmitt S, Jeltsch A. The human Dnmt2 has residual DNA-(cytosine-C5) methyltransferase activity. *J Biol Chem*. 2003;278:31717-21.
- Goll MG, Kirpekar F, Maggert KA, et al. Methylation of tRNA^{Asp} by the DNA methyltransferase homolog Dnmt2. *Science*. 2006;311:395-8.
- Deplus R, Brenner C, Burgers WA, et al. Dnmt3L is a transcriptional repressor that recruits histone deacetylase. *Nucleic Acids Res*. 2002;30:3831-8.
- Chedin F, Lieber MR, Hsieh CL. The DNA methyltransferase-like protein DNMT3L stimulates de novo methylation by Dnmt3a. *Proc Natl Acad Sci U S A*. 2002;99:16916-21.
- Hata K, Okano M, Lei H, Li E. Dnmt3L cooperates with the Dnmt3 family of de novo DNA methyltransferases to establish maternal imprints in mice. *Development*. 2002;129:1983-93.
- Brown SE, Fraga MF, Weaver IC, Berdasco M, Szyf M. Variations in DNA methylation patterns during the cell cycle of HeLa cells. *Epigenetics*. 2007;2:54-65.
- Bhattacharya SK, Ramchandani S, Cervoni N, Szyf M. A mammalian protein with specific demethylase activity for mCpG DNA. *Nature*. 1999;397:579-83.
- Watt F, Molloy PL. Cytosine methylation prevents binding to DNA of a HeLa cell transcription factor required for optimal expression of the adenovirus major late promoter. *Genes Dev*. 1988;2:1136-43.
- Drewell RA, Goddard CJ, Thomas JO, Surani MA. Methylation-dependent silencing at the H19 imprinting control region by MeCP2. *Nucleic Acids Res*. 2002;30:1139-44.
- Fujita N, Watanabe S, Ichimura T, et al. Methyl-CpG binding domain 1 (MBD1) interacts with the Suv39h1-HP1 heterochromatic complex for DNA methylation-based transcriptional repression. *J Biol Chem*. 2003;278:24132-8.
- Hoal-van Helden EG, van Helden PD. Age-related methylation changes in DNA may reflect the proliferative potential of organs. *Mutat Res*. 1989;219:263-6.
- Cooney CA, Dave AA, Wolff GL. Maternal methyl supplements in mice affect epigenetic variation and DNA methylation of offspring. *J Nutr*. 2002;132:2393S-400S.
- Salvador RA, Burton RM. Inhibition of the methylation of nicotinamide by chlorpromazine. *Biochem Pharmacol*. 1965;14:1185-96.
- Richardson BC, Liebling MR, Hudson JL. CD4+ cells treated with DNA methylation inhibitors induce autologous B cell differentiation. *Clin Immunol Immunopathol*. 1990;55:368-81.
- Schauenstein K, Rossi K, Csordas A. Differential inhibition of mitogen induced T cell proliferation by 5-azacytidine and cytosine-arabioside. *Biochem Biophys Res Commun*. 1988;151:548-53.
- Quddus J, Johnson KJ, Gavalcin J, et al. Treating activated CD4+ T cells with either of two distinct DNA methyltransferase inhibitors, 5-azacytidine or procainamide, is sufficient to cause a lupus-like disease in syngeneic mice. *J Clin Invest*. 1993;92:38-53.
- Lin SY, Hsieh SC, Lin YC, et al. A whole genome methylation analysis of systemic lupus erythematosus: hypomethylation of the IL10 and IL1R2

- promoters is associated with disease activity. *Genes Immun.* 2012;13:214–20.
33. Deng C, Kaplan MJ, Yang J, et al. Decreased Ras-mitogen-activated protein kinase signaling may cause DNA hypomethylation in T lymphocytes from lupus patients. *Arthritis Rheum.* 2001;44:397–407.
34. Sawalha AH, Jeffries M, Webb R, et al. Defective T-cell ERK signaling induces interferon-regulated gene expression and overexpression of methylation-sensitive genes similar to lupus patients. *Genes Immun.* 2008;9:368–78.
35. Marlin SD, Springer TA. Purified intercellular adhesion molecule-1 (ICAM-1) is a ligand for lymphocyte function-associated antigen 1 (LFA-1). *Cell.* 1987;51:813–9.
36. Lu Q, Ray D, Gutsch D, Richardson B. Effect of DNA methylation and chromatin structure on ITGAL expression. *Blood.* 2002;99:4503–8.
37. Richardson BC, Strahler JR, Pivrotto TS, et al. Phenotypic and functional similarities between 5-azacytidine-treated T cells and a T cell subset in patients with active systemic lupus erythematosus. *Arthritis Rheum.* 1992;35:647–62.
38. Kaplan MJ, Deng C, Yang J, Richardson BC. DNA methylation in the regulation of T cell LFA-1 expression. *Immunol Invest.* 2000;29:411–25.
39. Lu Q, Kaplan M, Ray D, et al. Demethylation of ITGAL (CD11a) regulatory sequences in systemic lupus erythematosus. *Arthritis Rheum.* 2002;46:1282–91.
40. Oelke K, Lu Q, Richardson D, et al. Overexpression of CD70 and overstimulation of IgG synthesis by lupus T cells and T cells treated with DNA methylation inhibitors. *Arthritis Rheum.* 2004;50:1850–60.
41. Lu Q, Wu A, Richardson BC. Demethylation of the same promoter sequence increases CD70 expression in lupus T cells and T cells treated with lupus-inducing drugs. *J Immunol.* 2005;174:6212–9.
42. Luo Y, Zhao M, Lu Q. Demethylation of promoter regulatory elements contributes to CD70 overexpression in CD4+ T cells from patients with subacute cutaneous lupus erythematosus. *Clin Exp Dermatol.* 2010;35:425–30.
43. Yin H, Zhao M, Wu X, et al. Hypomethylation and overexpression of CD70 (TNFSF7) in CD4+ T cells of patients with primary Sjogren's syndrome. *J Dermatol Sci.* 2010;59:198–203.
44. Jiang H, Xiao R, Lian X, et al. Demethylation of TNFSF7 contributes to CD70 overexpression in CD4+ T cells from patients with systemic sclerosis. *Clin Immunol.* 2012;143:39–44.
45. Ma L, Zhou Z, Wang H, et al. Increased expressions of DNA methyltransferases contribute to CD70 promoter hypomethylation and over expression of CD70 in ITP. *Mol Immunol.* 2011;48:1525–31.
46. Mosmann TR, Sad S. The expanding universe of T-cell subsets: Th1, Th2 and more. *Immunol Today.* 1996;17:138–46.
47. Yung RL, Qudus J, Chrisp CE, Johnson KJ, Richardson BC. Mechanism of drug-induced lupus. I. Cloned Th2 cells modified with DNA methylation inhibitors in vitro cause autoimmunity in vivo. *J Immunol.* 1995;154:3025–35.
48. Yano S, Ghosh P, Kusaba H, Buchholz M, Longo DL. Effect of promoter methylation on the regulation of IFN-gamma gene during in vitro differentiation of human peripheral blood T cells into a Th2 population. *J Immunol.* 2003;171:2510–6.
49. Katamura K, Fukui T, Kiyomasu T, et al. IL-4 and prostaglandin E2 inhibit hypomethylation of the 5' regulatory region of IFN-gamma gene during differentiation of naive CD4+ T cells. *Mol Immunol.* 1998;35:39–45.
50. Lee DU, Agarwal S, Rao A. Th2 lineage commitment and efficient IL-4 production involves extended demethylation of the IL-4 gene. *Immunity.* 2002;16:649–60.
51. Kwon NH, Kim JS, Lee JY, Oh MJ, Choi DC. DNA methylation and the expression of IL-4 and IFN-gamma promoter genes in patients with bronchial asthma. *J Clin Immunol.* 2008;28:139–46.
52. Mi XB, Zeng FQ. Hypomethylation of interleukin-4 and -6 promoters in T cells from systemic lupus erythematosus patients. *Acta Pharmacol Sin.* 2008;29:105–12.
53. Lal G, Zhang N, van der Touw W, Ding Y, et al. Epigenetic regulation of Foxp3 expression in regulatory T cells by DNA methylation. *J Immunol.* 2009;182:259–73.
54. Baron U, Floess S, Wieczorek G, et al. DNA demethylation in the human FOXP3 locus discriminates regulatory T cells from activated FOXP3(+) conventional T cells. *Eur J Immunol.* 2007;37:2378–89.
55. Kim HP, Leo. Abnormal DNA methylation in CD4+ T cells from people with latent autoimmune diabetes in adults. *nard WJ. CREB/ATF-dependent T cell receptor-induced FoxP3 gene expression: a role for DNA methylation. J Exp Med.* 2007;204:1543–51.
56. Li Y, Zhao M, Hou C, et al. Abnormal DNA methylation in CD4+ T cells from people with latent autoimmune diabetes in adults. *Diabetes Res Clin Pract.* 2011;94:242–8.
57. Podack ER, Hengartner H, Lichtenheld MG. A central role of perforin in cytotoxicity? *Annu Rev Immunol.* 1991;9:129–57.
58. Lu Q, Wu A, Ray D, et al. DNA methylation and chromatin structure regulate T cell perforin gene expression. *J Immunol.* 2003;170:5124–32.
59. Xiao R, Ding Y, Lu QJ, et al. Effects of 5-azaC on methylation pattern of the perforin promoter of the perforin gene in normal human T cells. *Zhong Nan Da Xue Xue Bao Yi Xue Ban.* 2006;31:843–7.
60. Kaplan MJ, Lu Q, Wu A, Attwood J, Richardson B. Demethylation of promoter regulatory elements contributes to perforin overexpression in CD4+ lupus T cells. *J Immunol.* 2004;172:3652–61.
61. Luo Y, Zhang X, Zhao M, Lu Q. DNA demethylation of the perforin promoter in CD4(+) T cells from patients with subacute cutaneous lupus erythematosus. *J Dermatol Sci.* 2009;56:33–6.
62. Armenante F, Merola M, Furia A, Palmieri M. Repression of the IL-6 gene is associated with hypermethylation. *Biochem Biophys Res Commun.* 1999;258:644–7.

63. Nile CJ, Read RC, Akil M, Duff GW, Wilson AG. Methylation status of a single CpG site in the IL6 promoter is related to IL6 messenger RNA levels and rheumatoid arthritis. *Arthritis Rheum.* 2008;58:2686-93.
64. Ishida K, Kobayashi T, Ito S, et al. Interleukin-6 gene promoter methylation in rheumatoid arthritis and chronic periodontitis. *J Periodontol.* 2012;83:917-25.
65. Hodge DR, Xiao W, Clausen PA, Heidecker G, Szyf M, Farrar WL. Interleukin-6 regulation of the human DNA methyltransferase (HDNMT) gene in human erythroleukemia cells. *J Biol Chem.* 2001;276:39508-11.
66. Bruniquel D, Schwartz RH. Selective, stable demethylation of the interleukin-2 gene enhances transcription by an active process. *Nat Immunol.* 2003;4:235-40.
67. Murayama A, Sakura K, Nakama M, et al. A specific CpG site demethylation in the human interleukin 2 gene promoter is an epigenetic memory. *Embo J.* 2006;25:1081-92.
68. Li Y, Ohms SJ, Shannon FM, Sun C, Fan JY. IL-2 and GM-CSF are regulated by DNA demethylation during activation of T cells, B cells and macrophages. *Biochem Biophys Res Commun.* 2012;419:748.
69. Hedrich CM, Rauen T, Tsokos GC. cAMP-responsive element modulator (CREM)alpha protein signaling mediates epigenetic remodeling of the human interleukin-2 gene: implications in systemic lupus erythematosus. *J Biol Chem.* 2011;286:43429-36.
70. Zhou Y, Yuan J, Pan Y, et al. T cell CD40LG gene expression and the production of IgG by autologous B cells in systemic lupus erythematosus. *Clin Immunol.* 2009;132:362-70.
71. Liao J, Liang G, Xie S, et al. CD40L demethylation in CD4(+) T cells from women with rheumatoid arthritis. *Clin Immunol.* 2012;145:13-8.
72. Lian X, Xiao R, Hu X, et al. DNA demethylation of CD40L in CD4+ T cells from women with systemic sclerosis: a possible explanation for female susceptibility. *Arthritis Rheum.* 2012;64:2338-45.
73. Lu Q, Wu A, Tesmer L, Ray D, Yousif N, Richardson B. Demethylation of CD40LG on the inactive X in T cells from women with lupus. *J Immunol.* 2007;179:6352-8.
74. Lleo A, Liao J, Invernizzi P, et al. Immunoglobulin M levels inversely correlate with CD40 ligand promoter methylation in patients with primary biliary cirrhosis. *Hepatology.* 2012;55:153-60.
75. Renaudineau Y, Hillion S, Saraux A, Mageed RA, Youinou P. An alternative exon 1 of the CD5 gene regulates CD5 expression in human B lymphocytes. *Blood.* 2005;106:2781-9.
76. Youinou P, Renaudineau Y. CD5 expression in B cells from patients with systemic lupus erythematosus. *Crit Rev Immunol.* 2011;31:31-42.
77. Garaud S, Le Dantec C, Jousse-Joulin S, et al. IL-6 modulates CD5 expression in B cells from patients with lupus by regulating DNA methylation. *J Immunol.* 2009;182:5623-32.
78. Garaud S, Le Dantec C, Berthou C, Lydyard PM, Youinou P, Renaudineau Y. Selection of the alternative exon 1 from the cd5 gene down-regulates membrane level of the protein in B lymphocytes. *J Immunol.* 2008;181:2010-18.
79. Lei W, Luo Y, Lei W, et al. Abnormal DNA methylation in CD4+ T cells from patients with systemic lupus erythematosus, systemic sclerosis, and dermatomyositis. *Scand J Rheumatol.* 2009;38:369-74.
80. Richardson B, Scheinbart L, Strahler J, Gross L, Hanash S, Johnson M. Evidence for impaired T cell DNA methylation in systemic lupus erythematosus and rheumatoid arthritis. *Arthritis Rheum.* 1990;33:1665-73.
81. Tao J, Yang M, Chen Z, et al. Decreased DNA methyltransferase 3A and 3B mRNA expression in peripheral blood mononuclear cells and increased plasma SAH concentration in adult patients with idiopathic thrombocytopenic purpura. *J Clin Immunol.* 2008;28:432-9.
82. Ogasawara H, Okada M, Kaneko H, Hishikawa T, Sekigawa I, Hashimoto H. Possible role of DNA hypomethylation in the induction of SLE: relationship to the transcription of human endogenous retroviruses. *Clin Exp Rheumatol.* 2003;21:733-8.
83. Zhao M, Tang J, Gao F, et al. Hypomethylation of IL10 and IL13 promoters in CD4+ T cells of patients with systemic lupus erythematosus. *J Biomed Biotechnol.* 2010;931018.
84. Xu J, Zhang G, Cheng Y, et al. Hypomethylation of the HTR1A promoter region and high expression of HTR1A in the peripheral blood lymphocytes of patients with systemic lupus erythematosus. *Lupus.* 2011;20:678-89.
85. Sunahori K, Juang YT, Kyttaris VC, Tsokos GC. Promoter hypomethylation results in increased expression of protein phosphatase 2A in T cells from patients with systemic lupus erythematosus. *J Immunol.* 2011;186:4508-17.
86. Javierre BM, Fernandez AF, Richter J, et al. Changes in the pattern of DNA methylation associate with twin discordance in systemic lupus erythematosus. *Genome Res.* 2010;20:170-9.
87. Basu D, Liu Y, Wu A, et al. Stimulatory and inhibitory killer Ig-like receptor molecules are expressed and functional on lupus T cells. *J Immunol.* 2009;183:3481-7.
88. Li Y, Zhao M, Yin H, et al. Overexpression of the growth arrest and DNA damage-induced 45alpha gene contributes to autoimmunity by promoting DNA demethylation in lupus T cells. *Arthritis Rheum.* 2010;62:1438-47.
89. Jeffries MA, Dozmorov M, Tang Y, Merrill JT, Wren JD, Sawalha AH. Genome-wide. DNA methylation patterns in CD4+ T cells from patients with systemic lupus erythematosus. *Epigenetics.* 2011;6:593-601.
90. Liu CC, Fang TJ, Ou TT, et al. Global DNA methylation, DNMT1, and MBD2 in patients with rheumatoid arthritis. *Immunol Lett.* 2011;135:96-9.
91. Karouzakis E, Rengel Y, Jungel A, et al. DNA methylation regulates the expression of CXCL12 in rheumatoid arthritis synovial fibroblasts. *Genes Immun.* 2011;12:643-52.

92. Fu LH, Cong B, Zhen YF, et al. Methylation status of the IL-10 gene promoter in the peripheral blood mononuclear cells of rheumatoid arthritis patients. *Yi Chuan*. 2007;29:1357-61.
93. Takami N, Osawa K, Miura Y, et al. Hypermethylated promoter region of DR3, the death receptor 3 gene, in rheumatoid arthritis synovial cells. *Arthritis Rheum*. 2006;54:779-87.
94. Han J, Park SG, Bae JB, et al. The characteristics of genome-wide DNA methylation in naive CD4+ T cells of patients with psoriasis or atopic dermatitis. *Biochem Biophys Res Commun*. 2012;422:157-63.
95. Zhang P, Su Y, Chen H, Zhao M, Lu Q. Abnormal DNA methylation in skin lesions and PBMCs of patients with psoriasis vulgaris. *J Dermatol Sci*. 2010;60:40-2.
96. Zhang K, Zhang R, Li X, Yin G, Niu X. Promoter methylation status of p15 and p21 genes in HPP-CFCs of bone marrow of patients with psoriasis. *Eur J Dermatol*. 2009;19:141-6.
97. Zhang K, Zhang R, Li X, Yin G, Niu X, Hou R. The mRNA expression and promoter methylation status of the p16 gene in colony-forming cells with high proliferative potential in patients with psoriasis. *Clin Exp Dermatol*. 2007;32:702-8.
98. Ruchusatsawat K, Wongpiyabovorn J, Shuangshoti S, Hirankarn N, Mutirangura A. SHP-1 promoter 2 methylation in normal epithelial tissues and demethylation in psoriasis. *J Mol Med (Berl)*. 2006;84:175-82.
99. Chen M, Chen ZQ, Cui PG, et al. The methylation pattern of p16INK4a gene promoter in psoriatic epidermis and its clinical significance. *Br J Dermatol*. 2008;158:987-93.
100. Mastronardi FG, Noor A, Wood DD, Paton T, Moscarello MA. Peptidyl argininedeiminase 2 CpG island in multiple sclerosis white matter is hypomethylated. *J Neurosci Res*. 2007;85:2006-16.
101. Kumagai C, Kalman B, Middleton FA, Vyshkina T, Massa PT. Increased promoter methylation of the immune regulatory gene SHP-1 in leukocytes of multiple sclerosis subjects. *J Neuroimmunol*. 2012;246:51-7.
102. Calabrese R, Zampieri M, Mechelli R, et al. Methylation-dependent PAD2 upregulation in multiple sclerosis peripheral blood. *Mult Scler*. 2012;18:299-304.
103. Hedrich CM, Rauen T. Epigenetic patterns in systemic sclerosis and their contribution to attenuated CD70 signaling cascades. *Clin Immunol*. 2012;143:1-3.
104. Fradin D, Le Fur S, Mille C, et al. Association of the CpG methylation pattern of the proximal insulin gene promoter with type 1 diabetes. *PLoS One*. 2012;7:e36278.
105. Saito S, Kato J, Hiraoka S, et al. DNA methylation of colon mucosa in ulcerative colitis patients: correlation with inflammatory status. *Inflamm Bowel Dis*. 2011;17:1955-65.
106. Tahara T, Shibata T, Nakamura M, et al. Effect of MDR1 gene promoter methylation in patients with ulcerative colitis. *Int J Mol Med*. 2009;23:521-7.
107. Hsieh CJ, Klump B, Holzmann K, Borchard F, Gregor M, Porschen R. Hypermethylation of the p16INK4a promoter in colectomy specimens of patients with long-standing and extensive ulcerative colitis. *Cancer Res*. 1998;58:3942-5.
108. Moriyama T, Matsumoto T, Nakamura S, et al. Hypermethylation of p14 (ARF) may be predictive of colitic cancer in patients with ulcerative colitis. *Dis Colon Rectum*. 2007;50:1384-92.
109. Arasaradnam RP, Khoo K, Bradburn M, Mathers JC, Kelly SB. DNA methylation of ESR-1 and N-33 in colorectal mucosa of patients with ulcerative colitis (UC). *Epigenetics*. 2010;5:422-6.
110. Tahara T, Shibata T, Nakamura M, et al. Promoter methylation of protease-activated receptor (PAR2) is associated with severe clinical phenotypes of ulcerative colitis (UC). *Clin Exp Med*. 2009;9:125-30.
111. Wang T, Wang Z, Yang R. Thrombopoietic growth factors in the treatment of immune thrombocytopenic purpura. *Crit Rev Oncol Hematol*. 2011;77:172-83.
112. Panitsas FP, Theodoropoulou M, Kouraklis A, et al. Adult chronic idiopathic thrombocytopenic purpura (ITP) is the manifestation of a type-1 polarized immune response. *Blood*. 2004;103:2645-7.
113. Semple JW, Milev Y, Cosgrave D, et al. Differences in serum cytokine levels in acute and chronic autoimmune thrombocytopenic purpura: relationship to platelet phenotype and antiplatelet T-cell reactivity. *Blood*. 1996;87:4245-54.
114. Olsson B, Andersson PO, Jernas M, et al. T-cell-mediated cytotoxicity toward platelets in chronic idiopathic thrombocytopenic purpura. *Nat Med*. 2003;9:1123-4.
115. Semple JW, Provan D, Garvey MB, Freedman J. Recent progress in understanding the pathogenesis of immune thrombocytopenia. *Curr Opin Hematol*. 2010;17:590-5.
116. Hauch TW, Rosse WF. Platelet-bound complement (C3) in immune thrombocytopenia. *Blood*. 1977;50:1129-36.
117. Melnyk S, Pogribna M, Pogribny IP, Yi P, James SJ. Measurement of plasma and intracellular S-adenosylmethionine and S-adenosylhomocysteine utilizing coulometric electrochemical detection: alterations with plasma homocysteine and pyridoxal 5'-phosphate concentrations. *Clin Chem*. 2000;46:265-72.
118. De Abreu R, Lambooy L, Stet E, Vogels-Mentink T, Van den Heuvel L. Thiopurine induced disturbance of DNA methylation in human malignant cells. *Adv Enzyme Regul*. 1995;35:251-63.
119. El-Shiekh EH, Bessa SS, Abdou SM, El-Refaey WA. Role of DNA methyltransferase 3A mRNA expression in Egyptian patients with idiopathic thrombocytopenic purpura. *Int J Lab Hematol*. 2012;34:369-76.
120. Hoffman DR, Cornatzer WE, Duerre JA. Relationship between tissue levels of S-adenosylmethionine, S-adenylhomocysteine, and transmethylation reactions. *Can J Biochem*. 1979;57:56-65.
121. Chen ZP, Gu DS, Zhou ZP, et al. Decreased expression of MBD2 and MBD4 gene and genomic-wide

- hypomethylation in patients with primary immune thrombocytopenia. *Hum Immunol.* 2011;72:486–91.
122. Hutchins AS, Artis D, Hendrich BD, Bird AP, Scott P, Reiner SL. Cutting edge: a critical role for gene silencing in preventing excessive type 1 immunity. *J Immunol.* 2005;175:5606–10.
123. Ma L, Zhou Z, Jia H, et al. Effects of CD70 and CD11a in immune thrombocytopenia patients. *J Clin Immunol.* 2011;31:632–42.
124. Zhao H, Du W, Wang D, et al. The expression of IFN- γ , IL-4, Foxp3 and perforin genes are not correlated with DNA methylation status in patients with immune thrombocytopenic purpura. *Platelets.* 2012;21:137–43.
125. Chen Z, Zhou Z, Chen X, et al. Single nucleotide polymorphism in DNMT3B promoter and the risk for idiopathic thrombocytopenic purpura in Chinese population. *J Clin Immunol.* 2008;28:399–404.
126. Shaheen IA, Abukhalil RE, Ali DK, Afifi RA. DNMT3B promoter polymorphism and risk of immune thrombocytopenic purpura in pediatric Egyptians. *Blood Coagul Fibrinolysis.* 2012;23:636–9.
127. Zhao H, Du W, Gu D, et al. DNMT3B 579G>T promoter polymorphism and the risk for idiopathic thrombocytopenic purpura in a Chinese population. *Acta Haematol.* 2009;122:31–5.
128. Hendrich B, Hardeland U, Ng HH, Jiricny J, Bird A. The thymine glycosylase MBD4 can bind to the product of deamination at methylated CpG sites. *Nature.* 1999;401:301–4.
129. Zhao H, Du W, Wang D, et al. Single nucleotide polymorphism in the methyl-CpG binding domain 4 gene and the risk for immune thrombocytopenic purpura in Chinese population. *Platelets.* 2010;21:132–6.